

Rehydrat® Tabs

agilan® complementary feed

To support digestion
highly soluble, effervescent, rich in energy and electrolytes

For calves, piglets and poultry

Composition

Lactose, Sodium bicarbonate, Sodium chloride, Potassium chloride

Analytical constituents

Crude protein	5.6 %
Crude fat	1.4 %
Crude fiber	0.5 %
Crude ash	17.4 %
Sodium	6.5 %
Potassium	1.3 %
Chloride	4.5 %
Lysine	0.0 %
Methionine	0.0 %

Content of additives per kg

Vitamins: Vitamin C (3a300) 10,000 mg

Recommended use

For oral rehydration in *calves, piglets and poultry*.

Rehydrat® Tabs quickly and reliably compensate electrolyte and body water losses due to an adequate mineral compositions and energy deficiencies due to an energy-rich lactose.

Feeding recommendation

Calves: 1 tab/liter of lukewarm water

should be given additionally to the usual milk meals as 1 - 3 snacks per day in a daily amount corresponding to the severity of the cases for independent intake.

The Rehydrat® solution should be used for the entire duration of an altered fecal consistency.

Fecal consistency	Additional liquid requirement (in % of b.w.)	40 kg calf	
		per day	waterings per day
Thin mushy	approx. 5 %	2 l	2 l / 1 x daily
Soupy	approx. 10 %	4 l	2 l / 2 x daily
Watery	15 (-20) %	6 (-8 l)	2 l / 3 x daily

Piglets: 1 tab/ liter of lukewarm water

The Rehydrat® solution is offered to the piglets for autonomous intake. It is favourable to prepare the solution fresh several times a day. This maintains the piglets' curiosity and prevents odor absorption from the environment.

Poultry: Dissolve 1 tab/ 100 liters of drinking water.



Rehydrat[®] Tabs

Feeding instruction

The solution should be freshly prepared with clean water prior to each use.

Rehydrat[®] Tabs should be administered at intervals to the ration of whole milk / milk replacer. Due to lactose Rehydrat[®] Tabs are very rich in energy with a low osmolarity. Lactose maintains the lactase activity so that it is possible to return to milk feeding without any problems.

This product must not be used as a sole drinker for calves and piglets.

Recommended feeding period: 1 - 7 days.

In case of risk of, during periods of or after digestive disturbance.

It is recommended that a veterinarian's opinion be sought before use or before extending the feeding period.

Storage

Shelf life guarantee is only valid for unopened packages, stored at a cool and dry place.

Keep the product out of the reach of children!

Date of revision of the text: December 2022